

## Contributors

- ANTHONY A. ALBANESE, *Nutritional Research Laboratory, St. Luke's Convalescent Hospital, Greenwich, Connecticut*
- JAMES B. ALLISON, *Bureau of Biological Research, and the Department of Physiology and Biochemistry, Rutgers, The State University, New Brunswick, New Jersey*
- H. J. ALMQUIST, *The Grange Company, Modesto, California*
- CHARLES H. BARROWS, JR., *Gerontology Branch, National Heart Institute, National Institutes of Health, P.H.S., D.H.E. & W., Bethesda, Maryland*
- CLARENCE P. BERG, *Department of Biochemistry, College of Medicine, State University of Iowa, Iowa City, Iowa*
- BACON F. CHOW, *The Baltimore City Hospitals and Department of Biochemistry, School of Hygiene and Public Health, The Johns Hopkins University, Baltimore, Maryland*
- DOUGLAS V. FROST, *Abbott Laboratories, North Chicago, Illinois*
- REGINALD A. HIGGONS, *Nutritional Research Laboratory, St. Luke's Convalescent Hospital, Greenwich, Connecticut*
- RUTH M. LEVERTON, *Institute of Home Economics, United States Department of Agriculture, Washington, D. C.*
- H. H. MITCHELL, *Division of Animal Nutrition, University of Illinois, Urbana, Illinois*
- BERNARD L. OSER, *Food and Drug Research Laboratories, Inc., Maspeth, New York*
- MARY S. PARSHLEY, *Department of Pathology, College of Physicians and Surgeons, Columbia University, New York*
- HANS R. ROSENBERG, *Nutrition Section, E. I. Du Pont de Nemours & Co., Stine Laboratory, Newark, Delaware*
- HENRY S. SIMMS, *Department of Pathology, College of Physicians and Surgeons, Columbia University, New York*
- PEARL SWANSON, *Department of Food and Nutrition, Iowa State College, Ames, Iowa*
- R. J. WILLIAMS, *The Clayton Foundation Biochemical Institute and the University of Texas, Austin, Texas*